

Black Hole in a Model with Dilaton and Monopole Fields II

E. Kyriakopoulos¹

Department of Physics
National Technical University
157 80 Zografou, Athens, GREECE

Abstract

We present an exact black hole solution in a model having besides gravity a dilaton and a monopole field, which is a generalization of a black hole solution we have found. The new solution, as the previous one, has three free parameters, one of which can be identified with the monopole charge, and another with the ADM mass. Its metric is asymptotically flat, has two horizon, irremovable singularity only at $r = 0$, and the dilaton field is singular only at $r = 0$. The dominant and the strong energy condition are satisfied outside and on the external horizon. According to a formulation of the no hair conjecture the solution is "hairy". Also a reformulation of the model with two monopole fields is given, which results in the appearance of an additional symmetry and therefore in the appearance of a conserved dilaton charge.

PACS number(s): 04.20.Jb, 04.70.Bw, 04.20.Dw

Keywords: Black hole, Dilaton, Monopole, No-hair theorem

According to a formulation of the no hair conjecture [1] : "We say that in a given theory there is black hole hair when the space-time metric and the configuration of the other fields of a stationary black hole solution are not completely specified by the conserved charges defined at asymptotic infinity". Previously [2] in a four-dimensional model having besides gravity a dilaton and a monopole field we found a solution which is "hairy" according to the above formulation. Other formulations of the no hair conjecture can also be found in the literature [3]-[6]. In this work we shall generalize our previous model. Also we shall reformulate the model such that an additional symmetry appears and therefore a conserved dilaton charge can be defined.

¹E-mail: kyriakop@central.ntua.gr, Tel: 00302107722980, Fax: 00302107722932

Consider the action

$$\int d^4x \sqrt{-g} L = \int d^4x \sqrt{-g} \left\{ R - \frac{1}{2} \partial_\mu \psi \partial^\mu \psi - f(\psi) F_{\mu\nu} F^{\mu\nu} \right\} \quad (1)$$

$$f(\psi) = g_1 e^{(c+\sqrt{c^2+1})\psi} + g_2 e^{(c-\sqrt{c^2+1})\psi} \quad (2)$$

where R is the Ricci scalar, ψ is a dilaton field, c , g_1 and g_2 are real constants and $F_{\mu\nu}$ is a pure monopole field

$$F = Q \sin\theta d\theta \wedge d\phi \quad (3)$$

with Q its magnetic charge. From this action we find the following equations of motion

$$(\partial^\rho \psi)_{;\rho} - \frac{df}{d\psi} F_{\mu\nu} F^{\mu\nu} = 0 \quad (4)$$

$$(f F^{\mu\nu})_{;\mu} = 0 \quad (5)$$

$$R_{\mu\nu} = \frac{1}{2} \partial_\mu \psi \partial_\nu \psi + 2f(F_{\mu\sigma} F_\nu{}^\sigma - \frac{1}{4} g_{\mu\nu} F_{\rho\sigma} F^{\rho\sigma}) \quad (6)$$

We want to find static spherically symmetric solutions of the above equations which are asymptotically flat and have regular horizon. We write the metric in the form [7]

$$ds^2 = -\lambda^2 dt^2 + \lambda^{-2} dr^2 + \xi^2 d\Omega \quad (7)$$

where λ and ξ are functions of r only and $d\Omega = d\theta^2 + \sin^2\theta d\phi^2$. From the above metric and Eq (3) we get $F_{\mu\nu} F^{\mu\nu} = \frac{2Q^2}{\xi^4}$, and we can prove that Eq (5) is satisfied. The dilaton Eq (4) for $\psi = \psi(r)$ takes the form

$$(\lambda^2 \xi^2 \psi')' = 2 \frac{df}{d\psi} Q^2 \xi^{-2} \quad (8)$$

where prime denotes differentiation with respect to r . The non-vanishing components of the Ricci tensor of the metric (7) are R_{00} , R_{11} , R_{22} and $R_{33} = \sin^2\theta R_{22}$, and for the first three components we get respectively from Eqs (6) the relations

$$(\lambda^2)'' + (\lambda^2)'(\xi^2)'\xi^{-2} = 2fQ^2\xi^{-4} \quad (9)$$

$$-(\lambda^2)''\lambda^{-2} - 2(\xi^2)''\xi^{-2} - (\lambda^2)'(\xi^2)'\lambda^{-2}\xi^{-2} + [(\xi^2)']^2\xi^{-4} = (\psi')^2 - 2fQ^2\lambda^{-2}\xi^{-4} \quad (10)$$

$$-[\lambda^2(\xi^2)]' + 2 = 2fQ^2\xi^{-2} \quad (11)$$

Eqs (8)-(11) form a system of four equations for the three unknowns λ^2 , ξ^2 and ψ . We found the following solution of this system

$$\lambda^2 = \frac{(r+A)(r+B)}{r(r+\alpha)} \left(\frac{r}{r+\alpha}\right)^{\frac{c}{\sqrt{c^2+1}}}, \quad \xi^2 = r(r+\alpha) \left(\frac{r+\alpha}{r}\right)^{\frac{c}{\sqrt{c^2+1}}} \quad (12)$$

$$e^\psi = e^{\psi_0} \left(1 + \frac{\alpha}{r}\right)^{\frac{1}{\sqrt{c^2+1}}} \quad (13)$$

where A , B , α and ψ_0 are integration constants, provided that the following relations are satisfied

$$g_1 = \frac{AB(\sqrt{c^2+1}-c)}{2Q^2\sqrt{c^2+1}} e^{-(c+\sqrt{c^2+1})\psi_0} \quad (14)$$

$$g_2 = \frac{(\alpha-A)(\alpha-B)(\sqrt{c^2+1}+c)}{2Q^2\sqrt{c^2+1}} e^{-(c-\sqrt{c^2+1})\psi_0} \quad (15)$$

From Eqs (7) and (12) we get

$$ds^2 = -\frac{(r+A)(r+B)}{r(r+\alpha)} \left(\frac{r}{r+\alpha}\right)^{\frac{c}{\sqrt{c^2+1}}} dt^2 + \frac{r(r+\alpha)}{(r+A)(r+B)} \left(\frac{r+\alpha}{r}\right)^{\frac{c}{\sqrt{c^2+1}}} dr^2 \\ + r(r+\alpha) \left(\frac{r+\alpha}{r}\right)^{\frac{c}{\sqrt{c^2+1}}} d\Omega \quad (16)$$

Our solution is given by Eqs (3) and (13)-(16).

From Eq. (16) we get asymptotically

$$-g_{00} = 1 - \frac{\alpha(1 + \frac{c}{\sqrt{c^2+1}}) - A - B}{r} + O(r^{-2}) \quad (17)$$

Therefore the solution is asymptotically flat and its Arnowitt-Deser-Misner (ADM) mass M is given by

$$2M = \alpha(1 + \frac{c}{\sqrt{c^2+1}}) - A - B \quad (18)$$

It is obvious that ψ_0 is the asymptotic value of ψ . Also for the choice

$$\alpha > 0, \quad A < 0, \quad B < 0, \quad (19)$$

we have

$$g_1 > 0, \quad g_2 > 0, \quad M > 0, \quad (20)$$

and ψ is singular only at $r = 0$. We shall make this choice for α , A and B .

The solution has the integration constants α , A , B , ψ_0 and Q , which for given g_1 , g_2 and c must satisfy Eqs (14) and (15). Therefore only three of them are independent. Introducing the ADM mass M by the relation (18) we can take M , Q and ψ_0 as independent parameters and express the parameters α , A and B of the dilaton field and the metric in terms of them. This means that the solution has arbitrary mass, arbitrary magnetic charge and an additional arbitrary parameter. Therefore it is a "hairy" solution according to the definition given in Ref. [1]. We should point out that the well known GHS-GM solution of D. Garfinkle, G. T. Horowitz and A. Strominger [7], found previously by G. W. Gibbons and by G. W. Gibbons and K. Maeda [8] is not a "hairy" solution according to this definition [9].

The metric coefficient g_{rr} is singular at $r = -A$ and $r = -B$ while the coefficient g_{tt} is singular at $r = 0$ but not at $r = -\alpha$, since $\alpha > 0$. However the singularities at $r = -A$ and $r = -B$ are coordinate singularities. Indeed the Ricci scalar R and the curvature scalar $R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}$ are given by

$$R = \frac{\alpha^2(r+A)(r+B)}{2(c^2+1)r^3(r+\alpha)^3} \left(\frac{r}{r+\alpha}\right)^{\frac{c}{\sqrt{c^2+1}}} \quad (21)$$

$$R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} = \frac{P(r, \alpha, A, B)}{4(c^2+1)^2r^6(r+\alpha)^6} \left(\frac{r}{r+\alpha}\right)^{\frac{2c}{\sqrt{c^2+1}}} \quad (22)$$

where $P(r, \alpha, A, B)$ is a complicated polynomial of r , α , A and B . Therefore R and $R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}$ are regular at $r = -A$ and $r = -B$, which means that only at $r = 0$ we have a real, an irremovable singularity.

If we make the Eddington-Finkelstein type transformation

$$dt = dt' \pm \frac{(A+B)r' + AB}{(r' + A)(r' + B)} \left(\frac{r' + \alpha}{r'}\right)^{1 + \frac{c}{\sqrt{1+c^2}}} dr', \quad r = r', \quad \theta = \theta', \quad \phi = \phi' \quad (23)$$

the metric ds^2 of Eq. (16) takes the regular at $r = -A$ and $r = -B$ form

$$\begin{aligned} ds^2 = & -\frac{(r' + A)(r' + B)}{r'(r' + \alpha)} \left(\frac{r'}{r' + \alpha}\right)^{\frac{c}{\sqrt{c^2+1}}} dt'^2 + \frac{r' + \alpha}{r'^3} \{r'^2 - (A+B)r' - AB\} \\ & \times \left(\frac{r' + \alpha}{r'}\right)^{\frac{c}{\sqrt{c^2+1}}} dr'^2 \mp \frac{2}{r'^2} \{(A+B)r' + AB\} dr' dt' + r'(r' + \alpha) \left(\frac{r' + \alpha}{r'}\right)^{\frac{c}{\sqrt{c^2+1}}} d\Omega \end{aligned} \quad (24)$$

From the above expression we find that the radial null directions, i.e. the directions for which $ds^2 = d\theta = d\phi = 0$, are determined by the relations

$$r' \left(\frac{r'}{r' + \alpha} \right)^{\frac{c}{\sqrt{c^2+1}}} dt' \pm (r' + \alpha) dr' = 0 \quad (25)$$

$$r'(r' + A)(r' + B) \left(\frac{r'}{r' + \alpha} \right)^{\frac{c}{\sqrt{c^2+1}}} dt' \mp (r' + \alpha) \{ r'^2 - (A + B)r' - AB \} dr' = 0 \quad (26)$$

The solution of the above equations give the intersections of the light cone with the $t' - r'$ plane. From the numerical solution of these Eqs we find that the solution with the upper sign is a black hole solution with two horizons at $r' = -A$ and at $r' = -B$, and the solution with the lower sign is a white hole solution.

The energy-momentum tensor $T_{\mu\nu}$ of our solution is given by

$$\begin{aligned} T_{\mu\nu} &= \partial_\mu \psi \partial_\nu \psi + 4f F_{\mu\rho} F_\nu{}^\rho - g_{\mu\nu} \left\{ \frac{1}{2} \partial_\rho \psi \partial^\rho \psi + f F_{\rho\sigma} F^{\rho\sigma} \right\} \\ &= \frac{\alpha^2}{(c^2 + 1)r^2(r + \alpha)^2} \delta_{\mu r} \delta_{\nu r} + \left\{ \frac{2AB}{r^2} \left(1 - \frac{c}{\sqrt{c^2 + 1}} \right) + \frac{2(\alpha - A)(\alpha - B)}{(r + \alpha)^2} \right. \\ &\quad \times \left(1 + \frac{c}{\sqrt{c^2 + 1}} \right) \left. \right\} (\delta_{\mu\theta} \delta_{\nu\theta} + \sin^2 \theta \delta_{\mu\phi} \delta_{\nu\phi}) - g_{\mu\nu} \left\{ \frac{\alpha^2(r + A)(r + B)}{2(c^2 + 1)r^3(r + \alpha)^3} \right. \\ &\quad \left. + \frac{AB}{r^3(r + \alpha)} \left(1 - \frac{c}{\sqrt{c^2 + 1}} \right) + \frac{(\alpha - A)(\alpha - B)}{r(r + \alpha)^3} \left(1 + \frac{c}{\sqrt{c^2 + 1}} \right) \right\} \left(\frac{r}{r + \alpha} \right)^{\frac{c}{\sqrt{c^2+1}}} \quad (27) \end{aligned}$$

Calculating the eigenvalues of $T_{\mu\nu}$ we can show that it satisfies the dominant as well as the strong energy condition outside and on the external horizon.

If $A < B < 0$ and $\alpha > 0$ the Hawking temperature T_H of our solution is [10]

$$T_H = \frac{|\lambda^{2'}(-A)|}{4\pi} = \frac{(B - A)(-A)^{-1 + \frac{c}{\sqrt{c^2+1}}} (\alpha - A)^{-1 - \frac{c}{\sqrt{c^2+1}}}}{4\pi} \quad (28)$$

The previous model can be reformulated such that the Lagrangian of the new model has an additional symmetry. To do that let us consider the action

$$\int d^4x \sqrt{-g} L = \int d^4x \sqrt{-g} \left\{ R - \frac{1}{2} \partial_\mu \psi \partial^\mu \psi - h(\psi) F_{\mu\nu} F^{\mu\nu} - k(\psi) G_{\mu\nu} G^{\mu\nu} \right\} \quad (29)$$

$$h(\psi) = g_1 e^{(c + \sqrt{c^2+1})\psi}, \quad k(\psi) = g_2 e^{(c - \sqrt{c^2+1})\psi} \quad (30)$$

where R is the Ricci scalar, ψ is a dilaton field, c , g_1 and g_2 are real constants and $F_{\mu\nu}$ and $G_{\mu\nu}$ are two pure monopole fields

$$F = Q_1 \sin\theta d\theta \wedge d\phi, \quad G = Q_2 \sin\theta d\theta \wedge d\phi \quad (31)$$

where Q_1 and Q_2 are their magnetic charges respectively. The above action for $g_1 = g_2 = 1$ and $c = 0$ is the part of the $SO(4)$ version of the $N = 4$, $d = 4$ supergravity action $I_{SO(4)}$ without the axion field [11]. Also the above action is invariant under the transformations

$$g_{\mu\nu} \rightarrow g_{\mu\nu}, \quad \psi \rightarrow \psi + b, \quad F_{\mu\nu} \rightarrow e^{-\left(\frac{c+\sqrt{c^2+1}}{2}\right)b} F_{\mu\nu}, \quad G_{\mu\nu} \rightarrow e^{-\left(\frac{c-\sqrt{c^2+1}}{2}\right)b} G_{\mu\nu} \quad (32)$$

where b is a constant. From this symmetry the conserved dilaton current

$$j_\mu = \partial_\mu \psi + 2(c + \sqrt{c^2 + 1})h(\psi)F_{\mu\nu}A^\nu + 2(c - \sqrt{c^2 + 1})k(\psi)G_{\mu\nu}B^\nu \quad (33)$$

is obtained and a conserved dilaton charge can be defined.

From the above action the following equations of motion

$$(\partial^\rho \psi)_{;\rho} - \frac{dh(\psi)}{d\psi} F_{\mu\nu} F^{\mu\nu} - \frac{dk(\psi)}{d\psi} G_{\mu\nu} G^{\mu\nu} = 0 \quad (34)$$

$$\{h(\psi)F^{\mu\nu}\}_{;\mu} = 0, \quad \{k(\psi)G^{\mu\nu}\}_{;\mu} = 0 \quad (35)$$

$$\begin{aligned} R_{\mu\nu} = & \frac{1}{2} \partial_\mu \psi \partial_\nu \psi + 2h(\psi)(F_{\mu\sigma} F_\nu{}^\sigma - \frac{1}{4} g_{\mu\nu} F_{\rho\sigma} F^{\rho\sigma}) \\ & + 2k(\psi)(G_{\mu\sigma} G_\nu{}^\sigma - \frac{1}{4} g_{\mu\nu} G_{\rho\sigma} G^{\rho\sigma}) \end{aligned} \quad (36)$$

are obtained. To get static spherically symmetric solutions of these equations which are asymptotically flat and have regular horizon we write the metric in the form of Eq. (7). Then we get from Eqs (31) $F_{\mu\nu} F^{\mu\nu} = \frac{2Q_1^2}{\xi^4}$ and $G_{\mu\nu} G^{\mu\nu} = \frac{2Q_2^2}{\xi^4}$, and we can prove that Eqs (35) are satisfied. The dilaton Eq. (34) takes the form

$$(\lambda^2 \xi^2 \psi')' = 2\left(\frac{dh}{d\psi} Q_1^2 + \frac{dk}{d\psi} Q_2^2\right) \xi^{-2} \quad (37)$$

where prime denotes differentiation with respect to r as before. The non-vanishing components of the Ricci tensor of the metric (7) are R_{00} , R_{11} , R_{22}

and $R_{33} = \sin^2\theta R_{22}$, and for the first three components we get respectively from Eqs (36) the relations

$$(\lambda^2)'' + (\lambda^2)'(\xi^2)'\xi^{-2} = 2(hQ_1^2 + kQ_2^2)\xi^{-4} \quad (38)$$

$$-(\lambda^2)''\lambda^{-2} - 2(\xi^2)''\xi^{-2} - (\lambda^2)'(\xi^2)'\lambda^{-2}\xi^{-2} + [(\xi^2)']^2\xi^{-4} = (\psi')^2 - 2(hQ_1^2 + kQ_2^2)\lambda^{-2}\xi^{-4} \quad (39)$$

$$-[\lambda^2(\xi^2)']' + 2 = 2(hQ_1^2 + kQ_2^2)\xi^{-2} \quad (40)$$

Eqs (37)-(40) form a system of four equations for the three unknowns λ^2 , ξ^2 and ψ . The expressions (12) and (13) with A , B , α and ψ_0 arbitrary constants, give again a solution of this system provided that the following relations are satisfied

$$g_1 = \frac{AB(\sqrt{c^2+1}-c)}{2Q_1^2\sqrt{c^2+1}}e^{-(c+\sqrt{c^2+1})\psi_0} \quad (41)$$

$$g_2 = \frac{(\alpha-A)(\alpha-B)(\sqrt{c^2+1}+c)}{2Q_2^2\sqrt{c^2+1}}e^{-(c-\sqrt{c^2+1})\psi_0} \quad (42)$$

Therefore in this model our solution is given by Eqs (13), (16), (31), (41) and (42). However in this case ψ_0 is not really a parameter because of the symmetry (32). For example using this symmetry we can make $\psi_0 = 0$. This means that, since the solution has the parameters A , B , α , Q_1 and Q_2 , which must satisfy Eqs (41) and (42), it has three free parameters. We can

take as such parameters the magnetic charges Q_1 and Q_2 and the mass M , which is given by Eq.(18). The conserved dilaton charge D of the solution is

$$D = \frac{1}{4\pi} \int_S J_\mu dS^\mu = -\frac{\alpha}{\sqrt{c^2+1}} \quad (43)$$

where the integral is over a two-sphere at spatial infinity. This charge can be expressed in terms of the free parameters Q_1 , Q_2 and M . Thus according to Ref. [1] the above solution is not "hairy", and according to Ref. [12] it has a hair of secondary type.

I am very grateful to A. Kehagias for many illuminating discussions and suggestions.

References

- [1] D. Nunez, H. Quevedo and D. Sudarsky, Phys. Rev. Lett. **76**, 571 (1996).
- [2] E. Kyriakopoulos submitted for publication in Int. J. Mod. Phys. D December 2006 Special Issue.
- [3] R. Ruffini and J. A. Wheeler, Phys. Today **24** (1),30 (1971).
- [4] J. D. Bekenstein, Phys. Rev. D **5**, 1239 (1972), **5**, 2403 (1972), **51** R6608 (1995), gr-gc/9808028.
- [5] S. Coleman, J. Preskill and F. Wilczek, Nucl. Phys. B **378**, 175 (1992).
- [6] P. Bizon, Acta Phys. Pol. B **22**, 877 (1994), gr-gc/9402016,
- [7] D. Garfinkle, G. T. Horowitz and A. Strominger, Phys. Rev. D **43**, 3140 (1991), **45**, 3888(E) (1992).
- [8] G. W. Gibbons Nucl. Phys. B **207**, 337 (1982), G. W. Gibbons and K. Maeda, Nucl. Phys. B **298**, 741 (1988).
- [9] D. Sudarsky and T. Zannias Phys. Rev. D **58** 087502 (1998).
- [10] G. T. Horowitz hep-th/9210119.
- [11] R. Kallosh, A. Linde, T. Ortin, A. Peet and A. Van Proeyen Phys. Rev. D **46**, 5278 (1992).
- [12] P. Kanti, N. E. Mavromatos, J. Rizos, K. Tamvakis and E. Winstanley, Phys. Rev. D **54**, 5049 (1996).